

Laboratory reconnections between plasma Jets and Tori: the PROTO-SPHERA Experiment

Franco Alladio, CR-ENEA Frascati



AGILE, 16th Science Workshop
ASI Headquarter, Roma, May 18 2018



Outline

- **PROTO-SPHERA has achieved the full Phase-1 current (10 kA) of Plasma Centerpost**
- **Surprise: the experiment now produces routinely Argon & Hydrogen Toroids**
- **DC voltage on Centerpost electrodes drives Toroidal Current (magnetic reconnections)**
- **In 2019: an intermediate Phase-1½ for moving to Phase-2 on sound bases**
- **End of 2021: Phase-2 experiment ready to produce plasma**

Magnetic confinement basics

Toroidal magnetic trap:
the inhomogeneous $\mathbf{B}$
separates charges



e.g. $e^- \rightarrow \text{up}$
 $\text{ions}^+ \rightarrow \text{down}$



$\mathbf{E}$ vertical field, through $\mathbf{E} \wedge \mathbf{B}$, ...ejects plasma

A poloidal $\mathbf{B}_\perp$ magnetic field is required to contrast particle drifts

→ toroidal current has to flow in an axisymmetric plasma



(magneto-hydrostatic equilibrium) $\vec{\nabla} p = \vec{j} \wedge \vec{B}$
flux (...also p & j) surfaces must be Tori

$$\vec{B} \cdot \vec{\nabla} p = 0$$



$$\vec{j} \cdot \vec{\nabla} p = 0$$

Magnetic confinement basics at work!

Switch-off of PROTO-SPHERA discharges containing a torus:


the brightness of the torus is the last to vanish,
both in the full spectrum of visible light

#1224, 10 kA Hydrogen



#1219, 10 kA Hydrogen

as well as observed through a Balmer-alpha filter






Tokamaks (1953)

Igor Yevgenyevich Tamm (1895-1971)

Dmitri Ivanovich Sakharov (1921-1989)



- The toroidal component of the magnetic field is produced by coils wound around the plasma torus
- The central solenoid induces the plasma current (*break-down*) and heats the plasma torus (ohmic heating, through plasma resistivity)
- Other poloidal field coils control the radial position of the plasma torus



Tokamak drive: AC currents, induction

Plasma confining current inside Torus:
induced and sustained by a transformer
whose current varies in time ...but there
are limits: the transformer will break
beyond a given current limit...


- The plasma ohmic drive in a Tokamak: **moving closed flux surfaces**, “feeding” the plasma from outside, they are dissipated while they move toward the magnetic axis
- In tokamaks this process is due to the transformer current change:
- **Tokamaks cannot have steady drive!**




The main idea of PROTO-SPHERA

- “Conventional Tokamak”: **magnetic surfaces of toroidal plasma** surround a “**Metal Centerpost**”

Toroidal geometry of Vacuum vessel



Electric current flow



- **PROTO-SPHERA**: **magnetic surfaces of toroidal plasma** surround a “**Plasma Centerpost**”; slim **metal external legs** return the current

Cylindrical geometry of Vacuum vessel

...but electrodes required inside vessel




Cylindrical vacuum vessel geometry has obvious advantages!


← easy to access & and to repair...

PROTO-SPHERA drive: DC currents, reconnections


- In front of the electrodes: open magnetic field lines,
- Open magnetic field lines are wound in a circular direction
- Magnetic reconnections convert open $\vec{B}$ lines into closed $\vec{B}, \vec{j}$ lines wrapped around the spherical torus



$\vec{E}$ of Tokamaks relies on **induction**
efficient but **not forever...**




$\vec{E}$ of PROTO-SPHERA relies on $\vec{v} \wedge \vec{B}$
associated with **magnetic reconnections**



SUSTAINMENT OF TOROIDAL CURRENT BY RECONNECTIONS OBTAINED in 2018 & IS FOREVER!

Reconnections of Jets and Tori a common occurrence in Nature!



Hydrodynamics examples

"Air-rings" in water



Beluga




Etna Volcano

"Smoke-rings"
in air

Plasma examples: Solar Flares Magnetic Reconnections, Coronal Heating

**PROTO-SPHERA (max 1 s plasma) aim was:
sustain Torus for 1 resistive time ~ 70 ms**

**PROTO-SPHERA has already sustained
a low current torus (5 kA) for $\frac{1}{2}$ sec**




The two Phases of PROTO-SPHERA

In **2002** at Frascati an **International Workshop** advised to build the machine in 2 steps:


- Phase-1: demonstrate Plasma Centerpost's (fear of anode arc anchoring...)
- Phase-2: machine completed such as to produce the Spherical Torus

PF coils casings built as floating,
charged by the plasma




Also upper\lower extensions
& lids are floating

Phase-1: 8 PFcoils
'Group B' shaping coils



Centerpost $I_e = 10 \text{ kA}$
...2018 Torus $I_{ST} = 5 \text{ kA}...$


Phase-2: 8 PF coils + 10 PF
'Group A' compressing coils




Centerpost $I_e = 60 \text{ kA}$
ST current $I_{ST} = 120 \div 240 \text{ kA}$

Ideal MHD stability of PROTO-SPHERA


Spheromak tilt instability is due to
dipole of containing field opposite to toroidal plasma current dipole



“Group A” PF coils (compression coils)
dipole moment opposes Plasma dipole
but “Group B” PF coils (shaping coils)
dipole moment is aligned to Plasma dipole



DISK-SHAPED ELECTRODE-FACING PLASMA GUARANTEES IDEAL MHD STABILITY



Cutting shorter & shorter
the plasma centerpost
PROTO-SPHERA at 120 kA
of toroidal current in ST
gets destabilized;
in absence of cutting
stability extends to 240 kA
of toroidal current in ST

PROTO-SPHERA Electrodes



May 2015 annular anode lowered on top of machine

Low voltage (100-350 V)
between electrodes

at high-field tokamak
plasma density $\sim \text{few} \bullet 10^{20} \text{ m}^{-3}$



Up: hollow gas-puffed anode





Down: 3000° K heated cathode

Phase-1 (present) cathode
(54 = 18 x 3 W emitters): aim 10 kA

Phase-2 cathode, 6 x Phase-1
(324 = 108 x 3 W emitters): aim 60 kA


“Caduceus”-like W emitting spirals
have now survived > 1500 cycles




A SPACE-THRUSTER forerunner?

abandon vacuum vessel toroidal geometry, move to cylindrical one
... → **natural expulsion of charged fusion products (Space Thruster)**

*Due to filamentary nature of B field
a fundamental mathematical difference appears:*




*a hairy torus
can be combed*




*a hairy sphere
cannot!*

From one of the “**tufts**” of the sphere (...not combed)
very high velocity ($\sim$ MeV) charged fusion products emerge

Possible future application as a Space Fusion Thruster...



Nozzle observed in PROTO-SPHERA experiment (2015)



PROTO-SPHERA Plasma Centerpost rotates & therefore avoids anode arc-anchoring



Anode arc-anchoring means that a plasma arc discharge is localized on an anode spot



#614, 8.5 kA Argon
break-down $V_e \sim 90$ V, steady $V_e \sim 200$ V



#977, 10 kA Hydrogen
break-down $V_e \sim 320$ V, steady $V_e \sim 200$ V

Plasma rotation in Tokamaks
always stabilizes the plasma

best way of injecting fuel is
from inboard (high field side)

(Pisa University & its PlasmaTech spin-off) line-averaged electron density on plasma equator:
in Argon Centerpost $\langle n_e \rangle \sim I_e$
at $I_e = 10$ kA, $\langle n_e \rangle \sim 4 \cdot 10^{20} \text{ m}^{-3}$
in Hydrogen Centerpost
at $I_e = 10$ kA, $\langle n_e \rangle \sim 1.5 \cdot 10^{20} \text{ m}^{-3}$


No Anode-Arc Anchoring: Electrostatic plasma effects!

$\vec{E} \wedge \vec{B}$ plasma rotation does the trick!


magnetic field is up\down symmetric

but

electrostatic field not up\down symmetric




Lines: magnetic field



Color contours: electrostatic potential, Arrows: E field

Insulating & transparent vessel being built for Phase-2

will substitute the Aluminum vacuum vessel with a Polymetacrylate (PMMA) transparent and insulating vessel (9.5 cm thick, 2m $\odot$, 1.7 m high), endowed with 20 ports adding 2 further SS rings on top & bottom of the experiment, keeping all internal components attached to the existing SS upper\lower lid and extension



Reynolds
POLYMER TECHNOLOGY, INC.
Building the Impossible




“Magnetic Boundary Conditions” changed

4 external PF coils added ...fed in series with the internal PF coils,
from April 2018 a further Super-Capacitor power
supply feeds some of the external PF coils


Hydrogen Centerpost
at full current =10 kA
seemed quite inclined
to eject a Torus
...an added vertical field
...was used as midwife!

Plasma fired after 0.75 s of PF current
to allow for skin current diffusion in Al vessel

#983, 10 kA Hydrogen




4 external PF coils




March 2018 Torus formed in Argon plasmas

In absence of Torus formation




With Torus formation




$I_e = 3$ kA

$I_e = 5$ kA


$I_e = 10$ kA



#1098, Argon



#1096, Argon




#1091, Argon

Torus forms at the expected current: $I_e \sim 8$ kA

$I_{\text{Torus}} = 5$ kA

The “gentle Divertor” of the (barely visible) Argon Torus



lower Divertor fan is dominant & slightly shifts down the Torus

The Argon Torus has been sustained for up to ½ sec




Argon Torus

break-down $V_e \sim 200$ V

steady $V_e \sim 220$ V

#1160, Argon, X-points fit $I_{ST}=5$ kA

April 2018 Torus formed in Hydrogen Plasma




evident at plasma breakdown



Hydrogen Torus
break-down
 $V_e \sim 360 \text{ V}$
steady $V_e \sim 320 \text{ V}$

#1202, Hydrogen

& ...after ¼ sec



at plasma switch-off


The “uncouth Divertor” of the Hydrogen Torus

The present divertor plate
is the lower polycarbonate diaphragm:
an unsuitable choice of material!

But when we put it inside PROTO-SPHERA
we did not expect to be able to form a Torus!


A new lower metal diaphragm being built now;
in 2019 PROTO-SPHERA will have 2 suitable plates

#1202, Hydrogen




Reconnections, Helical States & Double Tori

#1211
Hydrogen



#1225
Hydrogen



Vertical field in excess? ...Double Torus!

...or Helical States ...or lop-sided Tori!

The only occurrence you should never ever consider is plasma disruption: gone for good!



PROTO-SPHERA will not need additional heating...? ...magnetic reconnections heat the Solar Corona!

Phase-2 power injected into plasma $> 250 \text{ V} \bullet 60 \text{ kA} = 15 \text{ MW}$...how much into Torus?


~ 0 MW, $T = 10 \text{ eV}$
“cold” plasma:
divertor studies





$< 1 \text{ MW}$, $T = 100 \text{ eV}$
“lukewarm” plasma:
magnetic reconnection
studies at $S \sim 10^4$



a few MW, $T = 1 \text{ keV}$, “hot” plasma at $\beta \sim 1$, no disruptions
no additional heating & no current drive required,
same T as a Tokamak, but 1/100 of the cost!



Crab Nebula remnant



Hopf foliation

