

NAG Library Routine Document

F06QFF

Note: before using this routine, please read the Users' Note for your implementation to check the interpretation of ***bold italicised*** terms and other implementation-dependent details.

1 Purpose

F06QFF performs the matrix-copy operation

$$B \leftarrow A$$

where A and B are m by n real general or trapezoidal matrices.

2 Specification

```
SUBROUTINE F06QFF(MATRIX, M, N, A, LDA, B, LDB)
INTEGER          M, N, LDA, LDB
double precision A(LDA,*), B(LDB,*)
CHARACTER*1      MATRIX
```

3 Description

None.

4 References

None.

5 Parameters

- | | | |
|----|--|--------------|
| 1: | <p>MATRIX – CHARACTER*1</p> <p><i>On entry:</i> the matrix type.</p> <p>MATRIX = 'G'
General matrix.</p> <p>MATRIX = 'U'
Upper trapezoidal matrix (upper triangular if $m = n$).</p> <p>MATRIX = 'L'
Lower trapezoidal matrix (lower triangular if $m = n$).</p> <p><i>Constraint:</i> MATRIX = 'G', 'U' or 'L'.</p> | <i>Input</i> |
| 2: | <p>M – INTEGER</p> <p><i>On entry:</i> m, the number of rows of the matrices A and B.</p> <p><i>Constraint:</i> $M \geq 0$.</p> | <i>Input</i> |
| 3: | <p>N – INTEGER</p> <p><i>On entry:</i> n, the number of columns of the matrices A and B.</p> <p><i>Constraint:</i> $N \geq 0$.</p> | <i>Input</i> |

4: A(LDA,*) – *double precision* array *Input*

Note: the second dimension of the array A must be at least N.

On entry: the m by n general or trapezoidal matrix A .

If MATRIX = 'U', A is upper trapezoidal and the elements of the array below the diagonal are not referenced.

If MATRIX = 'L', A is lower trapezoidal and the elements of the array above the diagonal are not referenced.

5: LDA – INTEGER *Input*

On entry: the first dimension of the array A as declared in the (sub)program from which F06QFF is called.

Constraint: $LDA \geq \max(1, M)$.

6: B(LDB,*) – *double precision* array *Output*

Note: the second dimension of the array B must be at least N.

On exit: the m by n general or trapezoidal matrix B .

If MATRIX = 'U', B is upper trapezoidal and the elements of the array below the diagonal are not referenced.

If MATRIX = 'L', B is lower trapezoidal and the elements of the array above the diagonal are not referenced.

7: LDB – INTEGER *Input*

On entry: the first dimension of the array B as declared in the (sub)program from which F06QFF is called.

Constraint: $LDB \geq \max(1, M)$.

6 Error Indicators and Warnings

None.

7 Accuracy

Not applicable.

8 Further Comments

None.

9 Example

None.
