

Lecture 1

Shear Alfvén waves: continuous spectrum

Initial-value approach and time asymptotic response

Fulvio Zonca

<http://www.afs.enea.it/zonca>

ENEA C.R. Frascati, C.P. 65 - 00044 - Frascati, Italy.

Institute for Fusion Theory and Simulation, Zhejiang University, Hangzhou 310027, P.R.C.

April 27.th, 2017

Physics of Alfvén waves and energetic particles in fusion plasmas,

Linear physics and stability in tokamaks - Spring 2017

April 27 – May 12 2017, IFTS – ZJU, Hangzhou

Lecture Plan and Teaching Material

Abstract: This *Introduction to Physics of Alfvén waves and energetic particles in fusion plasmas* is devoted to *Linear physics and stability in tokamaks* and will be lectured by Prof. Fulvio Zonca in Spring 2017. These Lectures will cover with pedagogical style the first part of *Reviews of Modern Physics* **88**, 015008 (2016). Prerequisite of the course is a preliminary knowledge of general plasma physics and classical mechanics. General theoretical aspects are treated in Lectures (1 ÷ 6), which are complemented by corresponding Q&A Sessions.

Main areas that will be explored are:

(I) Fluid description of Alfvén waves: one dimensional plasmas

i) Shear Alfvén waves: continuous spectrum.

Initial-value approach and time asymptotic response. ([Lecture 1](#))

(II) Fluid description of Alfvén waves: two dimensional plasmas

- ii) Shear Alfvén waves in tokamaks. Mode structure decomposition and ballooning-mode representation.
Shear Alfvén continuous spectrum and frequency gaps. ([Lecture 2](#))
- iii) Low-frequency Alfvén and acoustic fluctuations.
Toroidal Alfvén Eigenmodes ([Lecture 3](#))

(III) Kinetic description of Alfvén waves in tokamaks

- iv) The linear gyrokinetic equation and the wave particle resonance condition in toroidal geometry. ([Lecture 4](#))
- v) Kinetic energy principle and variational approach.
Kinetic Alfvén waves. ([Lecture 5](#))
- vi) Energetic particle excitations of shear Alfvén instabilities in tokamaks.
([Lecture 6](#))

Lecture Notes: Available in electronic form. At the end of each lecture and during the Q&A time, a list of specific reading material is given explicitly. Should you have difficulty in finding literatures and papers, please contact me at fulvio.zonca@enea.it

Exercises and Research Projects: In the lecture notes, Exercises (E) of various difficulty levels are suggested. These are meant to be important part of the lectures themselves. Possible topics for Research Projects (RP) are also indicated, which require significant more in depth analysis and work.

Q&A Sessions: Devoted to detailed discussions and analysis of technical aspects. They are meant to be complementary to general lectures and are a necessary addendum for full appreciation of the material presented in the course.

Time and Location: Lectures (9am-11am) will be held at IFTS on April 27 and May 4-6-8-10-12, with corresponding Q&A Sessions on April 27-28 (4pm-5pm) and May 5-8-9-11-12 (3pm-5pm).

Review of ideal MHD

- Continuity equation: $\frac{d}{dt}\rho + \rho \nabla \cdot \mathbf{v} = 0$
- Momentum equation: $\rho \frac{d}{dt}\mathbf{v} = \frac{\mathbf{J} \times \mathbf{B}}{c} - \nabla p$
- Equation of state: $\frac{d}{dt}p + \gamma p \nabla \cdot \mathbf{v} = 0$
- Ohm's law: $\mathbf{E}' = \mathbf{E} + \frac{\mathbf{v} \times \mathbf{B}}{c} = 0$
- Faraday's law: $\nabla \times \mathbf{E} = -\frac{1}{c} \frac{\partial \mathbf{B}}{\partial t}$
- Ampère's law: $\nabla \times \mathbf{B} = \frac{4\pi}{c} \mathbf{J}$

$$\frac{d}{dt} = \frac{\partial}{\partial t} + \mathbf{v} \cdot \nabla$$

Waves in an infinite homogeneous magnetized plasma

- Uniform $\mathbf{B} = B_0 \mathbf{z}$ (no eq. current), $\mathbf{v}_0 = 0$, but finite ρ_0 and p_0 .
- Three purely oscillation modes are found: ($v_A^2 = B_0^2/(4\pi\rho_0)$, $v_S^2 = \gamma p_0/\rho_0$)
 - Shear Alfvén Wave (SAW): $\omega^2 = k_{\parallel}^2 v_A^2$ ($\rho_1 = p_1 = \nabla \cdot \mathbf{v}_1 = 0$)
 - Fast and Slow Magnetoacoustic Waves (FMW/SMW):

$$\omega^2 = (k^2/2)(v_A^2 + v_S^2) [1 \pm (1 - \alpha^2)^{1/2}]$$

$$(\alpha^2 = 4(k_{\parallel}^2/k^2)v_A^2 v_S^2/(v_A^2 + v_S^2)^2)$$

E: Derive the three branches above. Show that, for $\beta = v_S^2/v_A^2 \ll 1$, the FMW becomes the Compressional Alfvén Wave (CAW) with $\omega^2 = k^2 v_A^2$ and $4\pi p_1/(B_0 B_{1z}) = O(\beta)$; meanwhile the SMW becomes the Sound Wave (SW) with $\omega^2 = k_{\parallel}^2 v_S^2$ and $v_{1y}/v_{1z} = O(\beta)$.

E: Ideal MHD implies $\mathbf{E}_{\parallel} = 0$; yet it gives the SW! Solve the apparent paradox.

Fluid description of Alfvén waves: one dimensional plasmas

- To illustrate how plasma nonuniformities modify the spectral properties of SAW, let us consider the simple configuration of a 1D plasma slab confined in straight magnetic field [Chen 74, Goedbloed 84].
- The equilibrium quantities (plasma density, pressure and magnetic field) are assumed to vary only along the x direction ($\varrho_{m0} = \varrho_{m0}(x)$, $P_0 = P_0(x)$, $\mathbf{B}_0 = \mathbf{B}_0(x)$) and the equilibrium magnetic field is taken to be,

$$\mathbf{B}_0(x) = B_{0z}(x)\mathbf{e}_z.$$

- Here, we will adopt the well-known one-fluid ideal MHD description; which, we note, is valid for $|\omega/\omega_{ci}| \ll 1$, $|k_\perp \rho_i|^2 \ll 1$ and $m_e \ll m_i$.

E: demonstrate that one-fluid ideal MHD description is valid for $|\omega/\omega_{ci}| \ll 1$, $|k_\perp \rho_i|^2 \ll 1$ and $m_e \ll m_i$.

- Introducing the standard notation for the displacement vector $\delta \mathbf{u} \equiv \partial_t \delta \boldsymbol{\xi}$ and applying Faraday's and ideal Ohm's laws, such that

$$\delta \mathbf{B} = \nabla \times (\delta \boldsymbol{\xi} \times \mathbf{B}_0) ,$$

it is straightforward to derive the following linearized force balance equation

$$\rho_{m0} \frac{\partial^2}{\partial t^2} \delta \boldsymbol{\xi} = -\nabla \delta \tilde{p} + \frac{1}{4\pi} (\mathbf{B}_0 \cdot \nabla)^2 \delta \boldsymbol{\xi} - \frac{1}{4\pi} \mathbf{B}_0 (\mathbf{B}_0 \cdot \nabla) (\nabla \cdot \delta \boldsymbol{\xi}) .$$

- The total (kinetic plus magnetic) perturbed pressure

$$\delta \tilde{p} = \delta P + \frac{B_0 \delta B_{\parallel}}{4\pi}$$

corresponds to the eigenvector for the compressional Alfvén wave.

- Denoting by Γ the ratio of specific heats, from the equation of state, the perturbed plasma pressure can be obtained as

$$\delta P = -\delta \xi_x \frac{dP_0}{dx} - \Gamma P_0 (\nabla \cdot \delta \boldsymbol{\xi}) = \delta P_{\text{conv}} + \delta P_{\text{comp}} .$$

- Note that, by adopting a fluid description, we have suppressed important kinetic effects such as wave-particle interaction in the compressional component of δP , δP_{comp} (Lecture 4).
- Use the following Fourier representation for the generic perturbed quantity f :

$$f(\mathbf{r}, t) = \hat{f}(x) e^{-i(\omega t - k_y y - k_{\parallel} z)} .$$

- The force balance equation yields [Chen 74]

$$D_A \delta \xi_{\parallel} = \frac{4\pi}{B_0^2} i k_{\parallel} \delta \tilde{p} + i k_{\parallel} \left(i k_{\parallel} \delta \xi_{\parallel} + i k_y \delta \xi_y + \frac{d}{dx} \delta \xi_x \right),$$

$$D_A \delta \xi_y = \frac{4\pi}{B_0^2} i k_y \delta \tilde{p},$$

$$D_A \delta \xi_x = \frac{4\pi}{B_0^2} \frac{d}{dx} \delta \tilde{p},$$

- Here, we have denoted $v_A = (4\pi \rho_{m0})^{-1/2} B_0$ the Alfvén speed, $k_{\parallel} = k_z$ and $\delta \xi_{\parallel} = \delta \xi_z$; while the local SAW dispersion function is

$$D_A \equiv (\omega^2 / v_A^2) - k_{\parallel}^2.$$

- Applying Faraday's and Ohm's laws to $\delta B_{\parallel}$ in the expression of $\delta \tilde{p}$, one can readily write $\delta \tilde{p}$ in terms of the three components of $\delta \xi$,

$$\delta \tilde{p} = -i k_{\parallel} \Gamma P_0 \delta \xi_{\parallel} - \left(\Gamma P_0 + \frac{B_0^2}{4\pi} \right) \left(i k_y \delta \xi_y + \frac{d}{dx} \delta \xi_x \right).$$

- Substituting $\delta\xi_{\parallel}$ and noting the equation for $\delta\xi_y$, one obtains

$$\delta\xi_y(x) = \frac{i\bar{\alpha}k_y}{\bar{\alpha}k_y^2 - D_A} \frac{d\delta\xi_x}{dx},$$

where $\beta(x) \equiv 8\pi P_0(x)/B_0^2(x)$ and

$$\bar{\alpha}(x) \equiv 1 + \frac{\Gamma\beta\omega^2}{2\omega^2 - \Gamma\beta k_{\parallel}^2 v_A^2}.$$

E: Derive the above results step by step. What can you do next? What is the physical meaning of the equation that you are about to derive?

- Combining results, one finally arrives at the following wave equation in terms of $\delta\xi_x$ [Chen 74]:

$$\frac{d}{dx} \left(\frac{B_0^2 D_A \bar{\alpha}}{\bar{\alpha} k_y^2 - D_A} \frac{d\delta\xi_x}{dx} \right) - B_0^2 D_A \delta\xi_x = 0.$$

- This equation describes SAWs coupled with compressional Alfvén and ion-sound waves by equilibrium nonuniformities.

E: Show that this is indeed the case. Why can you say that the compressional Alfvén wave is described by this equation?

- We note that the obtained differential equation – and hence its solution ξ_x – is singular at the points where $B_0^2 D_A \bar{\alpha} = 0$.

E: Discuss what is the condition for a singular layer to appear in a differential equation. Apply this argument to the obtained differential equation.

The shear Alfvén continuous spectrum

- Singularity defines the following two continuous spectra.
One is the SAW continuum given by $D_A = 0$; *i.e.*, by

$$\omega^2 = \omega_A^2(x) \equiv k_{\parallel}^2 v_A^2(x).$$

The other is the ion-sound continuum given by $\bar{\alpha} = 0$; *i.e.*, with $v_S^2(x) = \Gamma P_0(x)/\varrho_{m0}(x)$ being the ion sound speed,

$$\omega^2 = \omega_S^2(x) \equiv \frac{v_S^2(x) k_{\parallel}^2}{1 + v_S^2(x)/v_A^2(x)}.$$

- Note that the spectra are continuous, since the wave frequency varies continuously in the non-uniformity direction x .

- To gain further insights into properties of SAW continuum, let us suppress the ion sound wave by assuming a cold plasma; that is $\Gamma\beta \rightarrow 0$ or $\bar{\alpha} \rightarrow 1$.
- Assume the plasma is confined by two perfectly conducting plates at $z = 0$ and $z = L$ [Chen 95].
- Our governing equation then describes shear and compressional Alfvén waves coupled by plasma non-uniformities; $\varrho_{m0}(x)$ in this case.
- In this limit, $\delta\xi_{\parallel} = 0$, and it is sufficient to consider the perpendicular components of the force balance equation

$$D_A \delta \boldsymbol{\xi}_{\perp} = \boldsymbol{\nabla}_{\perp} \left(\frac{\delta B_{\parallel}}{B_0} \right) .$$

- Here, $\delta \boldsymbol{\xi}_{\perp} = \delta \xi_y \mathbf{e}_y + \delta \xi_x \mathbf{e}_x$, $\boldsymbol{\nabla}_{\perp} = ik_y \mathbf{e}_y + \mathbf{e}_x \partial_x$ and meanwhile D_A is the local shear Alfvén operator

$$D_A \equiv \frac{\omega^2}{v_A^2} + \frac{\partial^2}{\partial z^2} .$$

- The compressional component of the magnetic field fluctuation is given by

$$\frac{\delta B_{\parallel}}{B_0} = -\nabla \cdot \delta \xi_{\perp}.$$

E: Justify the above expression using the results obtained so far.

- We now decompose all perturbed fields f in the complete orthonormal set of shear Alfvén eigenfunctions $\psi_{A\ell}(z|x)$

$$f(\mathbf{r}, t) = e^{-i(\omega t - k_y y)} \sum_{\ell=1}^{\infty} \hat{f}_{\ell}(x_0, x_1) \psi_{A\ell}(z|x_1),$$

where $\psi_{A\ell}(z|x)$, with boundary conditions $\psi_{A\ell} = 0$ at $z = 0$ and L , satisfies the differential equation ($\omega_{A\ell}^2(x)$ defines the local SAW frequency)

$$\left[\frac{\partial^2}{\partial z^2} + \frac{\omega_{A\ell}^2(x)}{v_A^2(x)} \right] \psi_{A\ell}(z|x) = 0.$$

- Perturbed field decomposition takes into account the existence of continuous frequency spectra, with a corresponding singular eigenfunction behavior, introducing “fast” x_0 (singular) and “slow” x_1 (equilibrium) radial variables.
- Using the following definition of the internal product

$$\langle \psi_{A\ell} | \psi_{A\ell'} \rangle \equiv \int_0^L v_A^2 \psi_{A\ell} \psi_{A\ell'} dz = \delta_{\ell\ell'} ,$$

with $k_\ell = \ell\pi/L$ and $\omega_{A\ell}^2(x) = k_\ell^2 v_A^2(x)$, one has

$$\psi_{A\ell}(z|x) = (2/Lv_A^2)^{1/2} \sin k_\ell z .$$

- We now exploit the existence of two spatial scales (x_0 and x_1) and solve the problem via asymptotic expansions of the fluctuating fields

$$\hat{f}_\ell = \hat{f}_\ell^{(0)} + \hat{f}_\ell^{(1)} + \dots ,$$

where $|\hat{f}_\ell^{(1)} / \hat{f}_\ell^{(0)}| \approx |\partial_{x_1} / \partial_{x_0}|$ etc.

- Noting that $|D_A| \ll k_y^2$ for a typical shear Alfvén wave (*i.e.*, $k_{\parallel}^2 \ll k_y^2$), the perpendicular force balance equation with the expression for $\delta B_{\parallel}$ may be combined and yield, at the lowest order,

$$\left[\frac{\partial}{\partial x_0} \epsilon_{A\ell} \frac{\partial}{\partial x_0} - k_y^2 \epsilon_{A\ell} \right] \delta \hat{\xi}_{x\ell}^{(0)} = 0,$$

where $\epsilon_{A\ell} = \omega^2 - \omega_{A\ell}^2(x)$.

E: Derive the equation above step by step.

- Solutions become singular at the radial positions $x_{R\ell}$ where the “local dispersion relation”, $\epsilon_{A\ell} = 0$ ($\omega^2 = \omega_{A\ell}^2(x_{R\ell})$), is satisfied; *i.e.*, at those positions where the shear Alfvén continuous spectrum is resonantly excited.
- A natural definition of the “fast” (singular) variable is $x_0 \equiv x - x_{R\ell}$, since $\epsilon_{A\ell} = \epsilon'_{A\ell}(x_1)x_0$.

- It is readily shown that solutions that may be written in the form

$$\delta \hat{\xi}_{x\ell}^{(0)} = \frac{C_\ell(x_1)}{\epsilon'_{A\ell}(x_1)} \ln(x_0) .$$

- Meanwhile, at the lowest order, $\nabla \cdot \delta \mathbf{\xi}_{\perp\ell}^{(0)} = 0$, from which we get

$$\delta \hat{\xi}_{y\ell}^{(0)} = \frac{i}{k_y} \partial_{x_0} \delta \hat{\xi}_{x\ell}^{(0)} = \frac{i C_\ell(x_1)}{k_y \epsilon'_{A\ell}(x_1)} \frac{1}{x_0} .$$

- Using this expression along with the y component of the perpendicular force balance, one can show that $\delta B_{\parallel}$ is given as

$$\delta B_{\parallel} = e^{-i(\omega t - k_y y)} \sum_{\ell=1}^{\infty} \delta \hat{b}_{\parallel\ell}(x_1) \psi_{A\ell}(z|x_1) .$$

- Here, the $\delta\hat{b}_{\parallel\ell}$ functions depend only on the slow variable and are directly related to the functions $C_\ell(x_1)$ introduced above:

$$\frac{\delta\hat{b}_{\parallel\ell}(x_1)}{B_0} = \frac{\epsilon'_{A\ell}(x_1)x_0}{ik_y v_A^2(x_1)} \delta\hat{\xi}_{y\ell}^{(0)} = \frac{C_\ell(x_1)}{k_y^2 v_A^2(x_1)}.$$

- This result means that, in a non-uniform plasma, shear and compressional Alfvén waves are coupled together and their coupling is the origin of the singular solutions corresponding to the “local” shear Alfvén oscillations of the continuous spectrum $\omega^2 = \omega_{A\ell}^2(x)$.
- The group velocity of shear Alfvén waves is directed along the magnetic field lines (z -direction in the present case), whereas the compressional wave generally carries energy across the field itself.
- Thus, CAW “piles up” wave energy at the radial location where the shear Alfvén spectrum is resonantly excited, explaining the origin of “local singular oscillations” [Chen 95].

Resonant excitation and resonant absorption

- The concept of resonant absorption of shear Alfvén waves is introduced by [Chen 1974]. In fact, a finite amount of wave energy can be absorbed at the resonant layer, $x_{R\ell}$.
- The time-averaged energy absorption rate, $d\langle W \rangle/dt$, is given by the Poynting energy flux into the infinitely narrow layer at $x_{R\ell}$. Thus [Chen 74], with L_y is the system extension in the y -direction,

$$\frac{d\langle W \rangle}{dt} = -\frac{cL_y}{8\pi} \operatorname{Re} \left\{ \int_0^L dz \int_{-\infty}^{\infty} \nabla \cdot \left[\left(\frac{i\omega}{c} \delta \boldsymbol{\xi} \times \mathbf{B}_0 \right) \times \delta \mathbf{B}^* \right] dx \right\} .$$

- Using the fluctuation decomposition and orthogonality relations, along with boundary conditions at $z = 0$ and $z = L$,

$$\frac{d\langle W \rangle}{dt} = -\frac{B_0 L_y}{8\pi} \operatorname{Re} \left\{ \sum_{\ell=1}^{\infty} \int_{-\infty}^{\infty} \frac{\partial}{\partial x} \left(-\frac{i\omega}{v_A^2} \delta \hat{\xi}_{x\ell} \delta \hat{b}_{\parallel\ell}^* \right) dx \right\} .$$

- The power absorption has contribution only at the resonant surface $x_{R\ell}$.
- Using the causality constraint $\text{Im } \omega \rightarrow 0^{(+)}$, it is possible to demonstrate that $\delta\hat{\xi}_{x\ell}^{(0)}$ (cf. p.18) has a jump given by

$$\Delta\delta\hat{\xi}_{x\ell}^{(0)} = -i\pi \operatorname{sgn} \left(\frac{\omega}{\epsilon'_{A\ell}(x_{R\ell})} \right) \frac{C_\ell(x_{R\ell})}{\epsilon'_{A\ell}(x_{R\ell})}.$$

- Thus, the power absorption becomes [Chen 95]

$$\begin{aligned} \frac{d\langle W \rangle}{dt} &= \frac{L_y}{8} k_y^2 \sum_{\ell=1}^{\infty} \frac{|\omega_{A\ell}(x_{R\ell})|}{|\epsilon'_{A\ell}(x_{R\ell})|} \left| \delta\hat{b}_{\parallel\ell}(x_{R\ell}) \right|^2 \\ &= \frac{L_y L}{8k_y^2} B_0^2 \sum_{\ell=1}^{\infty} |\omega'_{A\ell}(x_{R\ell})| \frac{\ell^2}{L^3 v_A^2(x_{R\ell})} \left| \Delta\delta\hat{\xi}_{x\ell}^{(0)} \right|^2. \end{aligned}$$

- This demonstrates resonant energy absorption at a rate $\propto \omega'_{A\ell}$. In turn, the resonant energy absorption of a given initial perturbation takes place on time scales $\approx (\omega'_{A\ell} \Delta x)^{-1}$, with Δx the perturbation *radial* extent.

Initial value approach

- The discussions on the singular solutions and resonant absorption are pertinent upon steady state excitations.
- It is also instructive to consider the wave perturbations from the initial-value approach.
- We are interested in the $|\omega_{\mathcal{A}l}t| \gg 1$ time asymptotic behaviors. Assuming, to be justified a posteriori, $|\partial_x| \gg |k_y|$, the governing equation (p. 17) with ω replaced by $i\partial_t$ then becomes

$$\frac{\partial}{\partial x} \left[\frac{\partial^2}{\partial t^2} + \omega_{\mathcal{A}l}^2(x) \right] \frac{\partial}{\partial x} \delta \hat{\xi}_{x\ell}(x, t) = 0.$$

- This equation can be straightforwardly integrated once and it yields

$$\frac{\partial}{\partial x} \delta \hat{\xi}_{x\ell}(x, t) = \hat{C}_\ell(x) e^{\pm i\omega_{\mathcal{A}l}(x)t},$$

- Here, $\hat{C}_\ell(x)$ is a function depending on equilibrium non-uniformities. Now, note that, as $\omega_{\mathcal{A}\ell}t \rightarrow \infty$ (the assumption $|\partial_x| \gg |k_y|$ is justified time asymptotically),

$$\partial_x \cong \pm i\omega'_{\mathcal{A}\ell}(x)t$$

and, thus,

$$\delta\hat{\xi}_{x\ell}(x, t) = \mp i \frac{\hat{C}_\ell(x)}{\omega'_{\mathcal{A}\ell}(x)t} e^{\pm i\omega_{\mathcal{A}\ell}(x)t}.$$


- From the condition $\nabla \cdot \delta\hat{\xi}_{\perp\ell} = 0$, one derives $\delta\hat{\xi}_{y\ell} = (i/k_y)\partial_x\delta\hat{\xi}_{x\ell}$; i.e.,

$$\delta\hat{\xi}_{y\ell}(x, t) = \frac{i}{k_y} \hat{C}_\ell(x) e^{\pm i\omega_{\mathcal{A}\ell}(x)t}.$$



- Insight in the dynamics associated with the resonant excitation of the shear Alfvén continuum and resonant wave absorption:

- the $\delta\hat{\xi}_{x\ell}$ component exhibits the characteristic $(1/t)$ decay via phase mixing of the continuous spectrum
- $\delta\hat{\xi}_{y\ell}$ shows undamped oscillations at frequencies corresponding to the shear Alfvén continuum


- These properties are nicely demonstrated by the satellite observations of magnetic field fluctuations in the Earth's magnetosphere [Engebretson 87]. B_R , B_E and B_N correspond to, respectively, $\delta B_x(\delta \hat{\xi}_x)$, $\delta B_y(\delta \hat{\xi}_y)$, $\delta B_{\parallel}$.



- The same behavior have also been demonstrated by ideal MHD initial value numerical simulations of Alfvén wave dynamics in a cylindrical plasma [Vlad 1999].
- Introducing (r, ϑ, z) as coordinate system in a cylindrical plasma of periodic length $2\pi R_0$ and radius a , $\delta \mathbf{u}(r, \vartheta, z, t) \equiv \partial_t \delta \boldsymbol{\xi} = e^{i(nz/R_0 - m\vartheta)} \delta \hat{\mathbf{u}}_{m,n}(r, t)$.



- The existence of “local singular oscillations” of the SAW continuous spectrum is further demonstrated by the Fourier frequency spectrum of $\delta\hat{u}_{1,1}^r(r, t)$ at several radial locations, reflecting the spatial structure of the continuum frequency $\omega_A(r/a)/\omega_A$ [Vlad 1999].



References and reading material

A. Hasegawa and L. Chen, Phys. Rev. Lett. **32**, 454 (1974).

L. Chen and A. Hasegawa, Phys. Fluids **17**, 1399 (1974).

L. Chen and A. Hasegawa, J. Geophys. Res. **79**, 1024 (1974).

J. P. Goedbloed, Physica **12D**, 107 (1984).

L. Chen and F. Zonca, Phys. Scr. **T60**, 81 (1995).

M. J. Engebretson, L. J. Zanetti, T. A. Potemra, W. Baumjohann, H. Lühr and M. H. Acuna, J. Geophys. Res. **92**, 10053 (1987).

G. Vlad, F. Zonca and S. Briguglio, Riv. Nuovo Cimento **22**, 1 (1999).